















Perineum: much distended and
 ascending surface of the stretched
 must, shows anterior wall of



80.

...mulerar några regle
 Rådfråga tillgängliga källor

Målningen »Eva«

Skolöförelse, Gårde - Bokundervisning



"...that death took root at the moment of birth
and man's only recourse thereafter was to water
and tend it"





ond half of pregnancy. A gain of about 1 pound (454 gm.) a week is suggested as a realistic maximum in the later months. Approximately half of all primigravidae from the twentieth to the thirtieth week will "naturally" gain this amount. More rigorous weight restrictions are not presently justified and may be harmful.

The Bones. The bones in pregnancy show increased vascularity, especially the red bone marrow. The spinal column is straightened; the lumbar curve is more prominent in front. The position of the woman's center of gravity tends to fall behind the pelvis for support. Therefore the shoulders back and the head. The curve of the back is exaggerated (Fig. 1). The pelvis is rotated slightly. These changes in the skeletal structure give the patient a peculiar attitude. However, the decided softening of the pelvic joints gives the patient a feeling of instability. These changes in the skeleton require a greater effort on the part of the mother to maintain an erect posture. An unaccustomed change in posture may cause backache by placing strains on certain muscles. The pendulous abdomens and changes in posture are ex-





6/31

Kate Brown







Handwritten notes in the bottom left corner, including "14/10/10" and "10/10/10".

Handwritten notes in the bottom center, including "10/10/10" and "10/10/10".

Handwritten notes in the bottom center, including "10/10/10" and "10/10/10".

Handwritten notes in the bottom right corner, including "10/10/10" and "10/10/10".



FIG. 1075.




Fig. 608



Fig. 609

FIG. 608. Uterus septum duplex or uterus bicornis. The vagina exists and there are two cervices with a longitudinal ridge between them. If the corpus also shows signs of splitting it is called uterus duplex bicornis (Fig. 609).

FIG. 609. Uterus duplex bicornis. The vagina exists and there are two cervices with a longitudinal ridge between them. If the corpus also shows signs of splitting it is called uterus duplex bicornis (Fig. 608).



pelvis and may be felt on bimanual examination as two cordlike, tender structures, coursing between the cervix and pelvic wall. Their walls are hypertrophic and hypotonic. This hypotonicity involves the uterus, the bowel, the ureters, the kidney pelvis and the abdominal wall. This general condition may be due to a disturbance in the balance of the autonomic and parasympathetic systems. The condition may be due to a disturbance in the balance of the autonomic and parasympathetic systems. The condition may be due to a disturbance in the balance of the autonomic and parasympathetic systems.



FIG. 180.





FIG. 180A. The bladder of the dog is held up by a continuous suture and the ureter is held up by a continuous suture.




FIG. 301. Unusual distribution of color in pregnancy. This patient also had large varicose veins on the vulva, in the vagina and in both lower extremities.

considerable
moved un-
Noseblee
nose is free
need be de
bleeding is



FIG. 302. Abruptio placentae. Considerable internal or external hemorrhage.

Internal or considerable



Admission of Guilt

























Sanna Claesson

@sanna_claesson

sanna.claesson@gmail.com

linkedin.com/in/sanna-claesson

UK + 44 7475 6165 12